

Author Index

Abraham, W., see Dragunow, M. (36) 349 Adan, R.A.H., Van der Kraan, M., Doornbos, R.P., Bär, P.R., Burbach, J.P.H. and Gispen, W.H.

Melanocortin receptors mediate α-MSH-induced stimulation of neurite outgrowth in Neuro 2A cells (36) 37

Afar, R., Silverman, R., Aguanno, A. and Albert, V.R.

Positive and negative elements contribute to the cell-specific expression of the rat dopamine β -hydroxylase gene (36) 79

Agnati, L.F., see Dasgupta, S. (36) 292 Aguanno, A., see Afar, R. (36) 79 Albert, V.R., see Afar, R. (36) 79 Arenas, E., see Dasgupta, S. (36) 292 Ashworth, R.G., see Margolis, R.L. (36) 219 Augood, S.J., see Harrington, K.A. (36) 157

Bär, P.R., see Adan, R.A.H. (36) 37 Barone, F.C., see Fan, L. (36) 287 Battenberg, E., see Morales, M. (36) 251 Becker, L.E., see Marks, A. (36) 343 Becker, W., Buttini, M., Limonta, S., Boddeke, H. and Joost, H.G. Distribution of the mRNA for protein phosphatase T in rat brain (36) 23 Bedell, M.A., see Nagasawa, M. (36) 1

Bjørås, M., Gjesdal, O., Erickson, J.D., Torp, R., Levy, L.M., Ottersen, O.P., Degree, M., Storm-Mathisen, J., Seeberg, E. and Danbolt, N.C.

Cloning and expression of a neuronal rat brain glutamate transporter (36) 163

Bleuel, Z., see Luque, J.M. (36) 357 Bloom, F.E., see Morales, M. (36) 251 Boddeke, H., see Becker, W. (36) 23 Born, W., see Eggenberger, M. (36) 127 Burbach, J.P.H., see Adan, R.A.H. (36) 37 Buttini, M., see Becker, W. (36) 23

Cambray-Deakin, M.A., see Przyborski, S.A. (36)179

Cansino, V.V., see Kohn, D.T. (36) 240 Cheetham, J.E., Martzen, M.R., Kazee, A.-M. and Coleman, P.D.

Gap-43 message levels in anterior cerebellum in Alzheimer's disease (36) 145

Chen, D.-Z., see Ohkuma, S. (36) 137 Chiaramello, A., see Haire, M.F. (36) 174 Chung, W.J., see Kindler, S. (36) 63

Cloëz-Tayarani, I., Wusher, N., Huerre, M. and Fillion, G.

Cellular localization of 5-HT_{1B} receptor mRNA in the rat olfactory tubercle: do GABAergic neurons express the 5-HT_{IR} gene? (36) 337

Coleman, P.D., see Cheetham, J.E. (36) 145 Combe, C., see Goujon, E. (36) 53 Copeland, N.G., see Nagasawa, M. (36) 1 Crawley, J.N., see Holmes, P.V. (36) 184 Curran, G.L., see Poduslo, J.F. (36) 280

Danbolt, N.C., see Bjørås, M. (36) 163 Dantzer, R., see Goujon, E. (36) 53 Dasgupta, S., Li, X.-M., Jansson, A., Finnman, U.-B., Matsui, T., Rinken, A., Arenas, E., Agnati, L.F. and Fuxe, K. Regulation of dopamine D2 receptor affinity by cholecystokinin octapeptide in fibroblast cells cotransfected with human CCK B and D₂L receptor cDNAs (36) 292

De Erausquin, G., see Isaacs, K.R. (36) 114 Degree, M., see Bjørås, M. (36) 163 De Lecea, L., see Morales, M. (36) 251 Dohanics, J., see Hoffman, G.E. (36) 201 Domowicz, M., see Li, H. (36) 309 Doornbos, R.P., see Adan, R.A.H. (36) 37 Dragunow, M., Abraham, W. and Hughes, P. Activation of NMDA and muscarinic receptors induces nur-77 mRNA in hippocampal neurons (36) 349 Dwyer, B.E., see Nishimura, R.N. (36) 227

Eggenberger, M., Flühmann, B., Muff, R., Lauber, M., Lichtensteiger, W., Hunziker, W., Fischer, J.A. and Born, W. Structure of a parathyroid hormone/parathyroid hormone-related peptide receptor of the human cerebellum and functional expression in human neuroblastoma SK-N-MC cells (36) 127

Ehlenfeldt, R., see Pavlou, O. (36) 268 Emson, P.C., see Harrington, K.A. (36) 157 Erickson, J.D., see Bjørås, M. (36) 163

Falkenberg, T., see Zachrisson, O. (36) 169 Fan, G., see Wick, M.J. (36) 322 Fan, L., Young, P.R., Barone, F.C., Feuerstein, G.Z., Smith, D.H. and McIntosh, T.K. Experimental brain injury induces differential expression of tumor necrosis factor- α mRNA in the CNS (36) 287

Feuerstein, G.Z., see Fan, L. (36) 287 Fillion, G., see Cloez-Tayarani, I. (36) 337 Finnman, U.-B., see Dasgupta, S. (36) 292

Fischer, J.A., see Eggenberger, M. (36) 127 Flühmann, B., see Eggenberger, M. (36) 127 Follesa, P., see Hu, X.-J. (36) 211 Foster, O.J.F., see Harrington, K.A. (36) 157 Furuyama, T., Inagaki, S., Iwahashi, Y., Wanaka, A. and Tohyama, M. Localization of mRNAs for Rlim-1, the rat Xlim-1 homolog, in the developing rat brain (36) 152

Fuxe, K., see Dasgupta, S. (36) 292

Garner, C.C., see Kindler, S. (36) 63 German, D.C., see Ng, M.C. (36) 329 Gispen, W.H., see Adan, R.A.H. (36) 37 Gjesdal, O., see Bjørås, M. (36) 163 Goujon, E., Parnet, P., Layé, S., Combe, C. and Dantzer, R. Adrenalectomy enhances pro-inflammatory cytokines gene expression, in the spleen, pituitary and brain of mice in response to lipopolysaccharide (36) 53

Grothe, C., see Meisinger, C. (36) 70 Grudt, T.J., Usowicz, M.M. and Henderson, G. Ca2+ entry following store depletion in SH-SY5Y neuroblastoma cells (36) 93

Guo, Y., see Kaplan, I.V. (36) 12

Haire, M.F. and Chiaramello, A. Transient expression of the basic helixloop-helix protein NSCL-2 in the mouse cerebellum during postnatal development

Hanbauer, I., see Isaacs, K.R. (36) 114 Harrington, K.A., Augood, S.J., Kingsbury, A.E., Foster, O.J.F. and Emson, P.C. Dopamine transporter (DAT) and synaptic vesicle amine transporter (VMAT2) gene expression in the substantia nigra of control and Parkinson's disease (36) 157

Hayashi, T., see Noga, M. (36) 261 Henderson, G., see Grudt, T.J. (36) 93 Hendrickson, A., see Luque, J.M. (36) 357 Hennig, A., see Li, H. (36) 309 Hertenstein, A., see Meisinger, C. (36) 70 Hoffman, G.E., Dohanics, J., Watson Jr., R.E. and Wiegand, S.J.

The hypothalamic ventromedial nucleus sends a met-enkephalin projection to the preoptic area's periventricular zone in the female rat (36) 201

Holmes, P.V. and Crawley, J.N. Olfactory bulbectomy increases preprogalanin mRNA levels in the rat locus coeruleus (36) 184

Horn, S., see Pavlou, O. (36) 268
Hu, X.-J., Follesa, P. and Ticku, M.K.
Chronic ethanol treatment produces a selective upregulation of the NMDA receptor subunit gene expression in mammalian cultured cortical neurons (36) 211
Huerre, M., see Cloëz-Tayarani, I. (36) 337
Hughes, P., see Dragunow, M. (36) 349

Hunziker, W., see Eggenberger, M. (36) 127

Iacopino, A.M., see Ng, M.C. (36) 329
Inagaki, S., see Furuyama, T. (36) 152
Isaacs, K.R., De Erausquin, G., Strauss, K.I.,
Jacobowitz, D.M. and Hanbauer, I.
Differential effects of excitatory amino acids on mesencephalic neurons expressing either calretinin or tyrosine hydroxylase in primary cultures (36) 114
Iwahashi, Y., see Furuyama, T. (36) 152

Jacobowitz, D.M., see Isaacs, K.R. (36) 114 Jansson, A., see Dasgupta, S. (36) 292 Jenkins, N.A., see Nagasawa, M. (36) 1 Joh, T.H., see Son, J.H. (36) 300 Joost, H.G., see Becker, W. (36) 23

Kaplan, I.V., Guo, Y. and Mower, G.D.
Immediate early gene expression in cat visual cortex during and after the critical period: differences between EGR-1 and Fos proteins (36) 12
Katsura, M., see Ohkuma, S. (36) 137
Kawakami, H., see Nakayama, T. (36) 189
Kazee, A.-M., see Cheetham, J.E. (36) 145
Kidwai, A.S., see Margolis, R.L. (36) 219
Kindler, S., Müller, R., Chung, W.J. and Garner, C.C.
Molecular characterization of dendritically localized transcripts encoding MAP2 (36) 63

Kingsbury, A.E., see Harrington, K.A. (36) 157
Kobayashi, S., see Kumai, T. (36) 197
Kohn, D.T., Tsai, K.-C., Cansino, V.V., Neve,
R.L. and Perrone-Bizzozero, N.I.
Role of highly conserved pyrimidine-rich
sequences in the 3' untranslated region of
the GAP-43 mRNA in mRNA stability and
RNA-protein interactions (36) 240

Kumai, T., Tanaka, M., Watanabe, M., Nakura, H., Tateishi, T. and Kobayashi, S. Elevated tyrosine hydroxylase mRNA levels in medulla oblongata of spontaneously hypertensive rats (36) 197

Kuriyama, K., see Ohkuma, S. (36) 137

Lamed, Y., see Paperna, T. (36) 101
Lauber, M., see Eggenberger, M. (36) 127
Layé, S., see Goujon, E. (36) 53
Lei, M., see Marks, A. (36) 343
Levy, L.M., see Bjørås, M. (36) 163
Li, H., Domowicz, M., Hennig, A. and
Schwartz, N.B.
S103L reactive chondroitin sulfate proteoglycan (aggrecan) mRNA expressed in developing chick brain and cartilage is encoded by a single gene (36) 309
Li, S.-H., see Margolis, R.L. (36) 219

Li, X.-M., see Dasgupta, S. (36) 292
Liang, C.-L., see Ng, M.C. (36) 329
Lichtensteiger, W., see Eggenberger, M. (36) 127
Limonta, S., see Becker, W. (36) 23
Lindefors, N., see Zachrisson, O. (36) 169
Loh, H.H., see Wick, M.J. (36) 322
Luque, J.M., Bleuel, Z., Hendrickson, A. and Richards, J.G.
Detection of MAO-A and MAO-B mRNAs in monkey brainstem by cross-hybridization

with human oligonucleotide probes (36) 357

Marches, F., see Ng, M.C. (36) 329 Margolis, R.L., Li, S.-H., Young, W.S., Wagster, M.V., Stine, O.C., Kidwai, A.S., Ashworth, R.G. and Ross, C.A. DRPLA gene (Atrophin-1) sequence and mRNA expression in human brain (36) 219 Marks, A., O'Hanlon, D., Lei, M., Percy, M.E. and Becker, L.E. Accumulation of \$100\beta mRNA and protein in cerebellum during infancy in Down syndrome and control subjects (36) 343 Martzen, M.R., see Cheetham, J.E. (36) 145 Matsui, T., see Dasgupta, S. (36) 292 McIntosh, T.K., see Fan, L. (36) 287 Meisinger, C., Hertenstein, A. and Grothe, C. Fibroblast growth factor receptor 1 in the adrenal gland and PC12 cells: developmental expression and regulation by extrinsic molecules (36) 70 Min, N., see Son, J.H. (36) 300 Mishina, M., see Nagasawa, M. (36) 1 Morales, M., Battenberg, E., De Lecea, L., Sanna, P.P. and Bloom, F.E.

Nagasawa, M., Sakimura, K., Mori, K.J., Bedell, M.A., Copeland, N.G., Jenkins, N.A. and Mishina, M.

Gene structure and chromosomal localization of the mouse NMDA receptor channel

Cellular and subcellular immunolocalization

of the type 3 serotonin receptor in the rat

central nervous system (36) 251

Mower, G.D., see Kaplan, I.V. (36) 12

Muff, R., see Eggenberger, M. (36) 127

Mori, K.J., see Nagasawa, M. (36) 1

Müller, R., see Kindler, S. (36) 63

subunits (36) 1
Nakamura, S., see Nakayama, T. (36) 189
Nakayama, T., Kawakami, H., Tanaka, K. and
Nakamura, S.
Expression of three glutamate transporter
subtype mRNAs in human brain regions

and peripheral tissues (36) 189 Nakura, H., see Kumai, T. (36) 197 Narihara, H., see Ohkuma, S. (36) 137 Neve, R.L., see Kohn, D.T. (36) 240

Ng, M.C., Iacopino, A.M., Quintero, E.M.,
 Marches, F., Sonsalla, P.K., Liang, C.-L.,
 Speciale, S.G. and German, D.C.
 The neurotoxin MPTP increases calbindin D_{28k} levels in mouse midbrain dopaminer gic neurons (36) 329

Nishimura, R.N. and Dwyer, B.E. Evidence for different mechanisms of induction of HSP70i: a comparison of cultured rat cortical neurons with astrocytes (36) 227

Noga, M. and Hayashi, T.

Ubiquitin gene expression following transient forebrain ischemia (36) 261

O'Hanlon, D., see Marks, A. (36) 343
Ohkuma, S., Katsura, M., Chen, D.-Z., Narihara, H. and Kuriyama, K.
Nitric oxide-evoked [³H]y-aminobutyric acid release is mediated by two distinct release mechanisms (36) 137

Orr, H.T., see Pavlou, O. (36) 268 Ottersen, O.P., see Bjørås, M. (36) 163

Paperna, T., Lamed, Y. and Teichberg, V.I.

CDNA cloning of chick brain α-amino-3hydroxy-5-methyl-4-isoxazolepropionic acid
receptors reveals conservation of structure,
function and post-transcriptional processes
with mammalian receptors (36) 101

Parnet, P., see Goujon, E. (36) 53
Pavlou, O., Ehlenfeldt, R., Horn, S. and Orr, H.T.
Isolation, characterization and in vivo analysis of the murine calbindin-D_{28K} upstream regulatory region (36) 268

Percy, M.E., see Marks, A. (36) 343 Perrone-Bizzozero, N.I., see Kohn, D.T. (36) 240

Poduslo, J.F. and Curran, G.L.

Permeability at the blood-brain and bloodnerve barriers of the neurotrophic factors:
NGF, CNTF, NT-3, BDNF (36) 280

Przyborski, S.A. and Cambray-Deakin, M.A. Developmental regulation of α-tubulin mR-NAs during the differentiation of cultured cerebellar granule cells (36) 179

Puchacz, E., Stumpf, W.E., Stachowiak, E.K. and Stachowiak, M.K.
Vitamin D increases expression of the tyrosine hydroxylase gene in adrenal medullary cells (36) 193

Quintero, E.M., see Ng, M.C. (36) 329

Richards, J.G., see Luque, J.M. (36) 357 Rinken, A., see Dasgupta, S. (36) 292 Ross, C.A., see Margolis, R.L. (36) 219

Sakimura, K., see Nagasawa, M. (36) 1
Sanna, P.P., see Morales, M. (36) 251
Schwartz, N.B., see Li, H. (36) 309
Seeberg, E., see Bjørås, M. (36) 163
Silverman, R., see Afar, R. (36) 79
Smith, D.H., see Fan, L. (36) 287
Son, J.H., Min, N. and Joh, T.H.
Early ontogeny of catecholaminergic cell lineage in brain and peripheral neurons monitored by tyrosine hydroxylase-lacZ transgene (36) 300
Sonsalla, P.K., see Ng, M.C. (36) 329
Speciale, S.G., see Ng, M.C. (36) 329
Stachowiak, F.K., see Puchacz, F. (36) 193

Speciale, S.G., see Ng, M.C. (36) 329
Stachowiak, E.K., see Puchacz, E. (36) 193
Stachowiak, M.K., see Puchacz, E. (36) 193
Stine, O.C., see Margolis, R.L. (36) 219
Storm-Mathisen, J., see Bjørås, M. (36) 163
Strauss, K.I., see Isaacs, K.R. (36) 114
Stumpf, W.E., see Puchacz, E. (36) 193

Takagi, T., see Taniguchi, H. (36) 45
Tanaka, K., see Nakayama, T. (36) 189
Tanaka, M., see Kumai, T. (36) 197
Taniguchi, H., Tohyama, M. and Takagi, T.
Cloning and expression of a novel gene for a protein with leucine-rich repeats in the developing mouse nervous system (36) 45
Tateishi, T., see Kumai, T. (36) 197
Teichberg, V.I., see Paperna, T. (36) 101
Ticku, M.K., see Hu, X.-J. (36) 211
Tohyama, M., see Furuyama, T. (36) 152
Tohyama, M., see Taniguchi, H. (36) 45
Torp, R., see Bjørås, M. (36) 163
Tsai, K.-C., see Kohn, D.T. (36) 240

Usowicz, M.M., see Grudt, T.J. (36) 93

Van der Kraan, M., see Adan, R.A.H. (36) 37

Wagster, M.V., see Margolis, R.L. (36) 219

Wanaka, A., see Furuyama, T. (36) 152

Wang, H.-L. and Wu, T.

 $G_{\alpha q/11}$ mediates neurotensin excitation of substantia nigra dopaminergic neurons (36)

Watson Jr., R.E., see Hoffman, G.E. (36) 201 Wick, M.J., Fan, G. and Loh, H.H. Expression of OBCAM-related cDNA clones in Cos 1 cells: evidence for a phos-

Watanabe, M., see Kumai, T. (36) 197

phatidylinositol linkage to the cell membrane (36) 322
Wiegand, S.J., see Hoffman, G.E. (36) 201
Wu, T., see Wang, H.-L. (36) 29
Wusher, N., see Cloëz-Tayarani, I. (36) 337

Young, P.R., see Fan, L. (36) 287 Young, W.S., see Margolis, R.L. (36) 219

Zachrisson, O., Falkenberg, T. and Lindefors, N.
Neuronal coexistence of trkB and glutamic acid decarboxylase₆₇ mRNAs in rat hip-pocampus (36) 169

